



Original Research Article

# Surgical Innovations and Outcomes in the Management of Rectal Cancer: A Departmental Study on Advanced Techniques and Postoperative Care

Dr. Md. Shahidul Islam<sup>1\*</sup>, Dr. kazi Shah Md. Abdullah<sup>2</sup>, Dr. Choudhury Md. Anwar Sadat<sup>3</sup>, Dr. Md. Imrul Islam<sup>4</sup>

<sup>1</sup>Consultant, Department of Surgery, Upazila Health Complex, Singra, Natore, Rajshahi

<sup>2</sup>Junior Consultant, Department of Surgery, 100 Bed Sadar Hospital, Shariatpur

<sup>3</sup>Assistant Professor Department of Surgery, Parkview Medical College and Hospital, Sylhet

<sup>4</sup>Senior consultant, Anaesthesia, 250 bed district hospital, Chapa Nawabganj, Rajshahi



**Citation:**

Md. Shahidul Islam, kazi Shah Md. Abdullah, Choudhury Md. Anwar Sadat, Md. Imrul Islam (2024). Surgical Innovations and Outcomes in the Management of Rectal Cancer: A Departmental Study on Advanced Techniques and Postoperative Care. *Asia Pac J Cancer Res*, 1(1), 14-22.

Received: 10/06/2024

Accepted: 22/07/2024

Published: 31/08/2024



**Copyright © 2024 The Author(s):**

This is an open-access article distributed under the terms of the Creative Commons Attribution 4.0 International License (CC BY-NC 4.0) which permits unrestricted use, distribution, and reproduction in any medium for non-commercial use provided the original author and source are credited.

**Abstract: Background:** Rectal cancer is a significant global health issue, accounting for a substantial proportion of colorectal cancer cases. Surgical innovations and comprehensive postoperative care have significantly improved patient outcomes. This study evaluates the impact of advanced surgical techniques and Enhanced Recovery After Surgery (ERAS) protocols on managing rectal cancer across multiple centers in the Rajshahi Division. **Objective:** The study aimed to assess the effectiveness of minimally invasive surgical techniques, including laparoscopic, -assisted, and transanal total mesorectal excision (taTME), compared to traditional open Surgery. It also examined the role of ERAS protocols in enhancing recovery and reducing complications. **Method:** This multicenter, retrospective study included 88 patients diagnosed with rectal cancer who underwent surgical treatment from June 2021 to June 2023 across various hospitals in the Rajshahi Division. Patients were categorized based on the type of Surgery received, and postoperative outcomes, including complications, hospital stay, and survival rates, were analyzed. Data were statistically evaluated using SPSS version 26, with significance at  $p < 0.05$ . **Result:** Of the 88 patients, 50% underwent laparoscopic Surgery, 20% -assisted surgery, 10% taTME, and 20% open surgery. Patients who underwent minimally invasive techniques had significantly shorter hospital stays (mean of 6 days) than open Surgery (mean of 10 days,  $p = 0.001$ ). The overall complication rate was lower in the minimally invasive group (17%) compared to the open surgery group (30%). Two-year survival rates were highest in the -assisted and taTME groups (90% and 92%, respectively). **Conclusions:** Advanced surgical techniques, particularly -assisted Surgery, and taTME, combined with ERAS protocols, significantly improve patient outcomes in rectal cancer management.

**Keywords:** Rectal cancer, Minimally invasive Surgery, taTME, ERAS protocols.

**Significance:** Minimally invasive techniques and ERAS protocols significantly enhance rectal cancer outcomes, reducing complications and improving survival and recovery.

## INTRODUCTION

Rectal cancer remains a significant global health concern, accounting for a substantial proportion of colorectal cancer cases [1]. With advancements in surgical techniques and postoperative care, there have been remarkable

improvements in patient outcomes, particularly in terms of survival rates, quality of life, and reduced morbidity. This study focuses on the surgical innovations and outcomes in managing rectal cancer within our department, emphasizing the adoption of advanced techniques and comprehensive postoperative care protocols. Rectal

cancer is a major public health issue worldwide, with its incidence and mortality rates varying significantly across different regions. According to the Global Cancer Observatory, rectal cancer accounted for approximately 30% of colorectal cancer cases in 2020, with over 700,000 new cases reported globally [2]. The rising incidence, particularly in low- and middle-income countries, underscores the need for innovative surgical approaches and improved postoperative care to enhance patient outcomes.

The management of rectal cancer has evolved significantly over the past few decades. Traditionally, Surgery has been the cornerstone of curative treatment, with total mesorectal excision (TME) being the gold standard [3]. However, the complexity of rectal anatomy and the proximity of the tumor to critical structures such as the pelvic nerves and sphincter muscles make Surgery challenging. Consequently, there has been a growing interest in developing advanced surgical techniques to improve resection margins, preserve organ function, and reduce postoperative complications. One of the most significant advancements in rectal cancer surgery is the adoption of minimally invasive techniques, such as laparoscopic and -assisted Surgery. These techniques have gained popularity due to their potential to reduce operative trauma, shorten hospital stays, and accelerate patient recovery. A study by Jiang *et al.*, demonstrated that laparoscopic Surgery for rectal cancer is associated with similar oncological outcomes compared to open Surgery, with the added benefit of reduced postoperative pain and quicker return to normal activities [4]. Similarly, -assisted Surgery, with its enhanced precision and skill, has been shown to improve the quality of TME and reduce the risk of positive circumferential resection margins [5].

Another innovative approach in rectal cancer surgery is the transanal total mesorectal excision (taTME), which offers a clear view of the distal rectum and the ability to perform a more precise dissection [6]. This technique is particularly beneficial in patients with mid to low-rectal tumors, where traditional TME may be challenging. Early results from studies comparing taTME with conventional TME have shown promising outcomes, including lower rates of conversion to

open Surgery, reduced intraoperative blood loss, and better preservation of bowel function [7]. Implementing these advanced surgical techniques in our department has been driven by improving patient outcomes, particularly in terms of sphincter preservation, reduced morbidity, and enhanced recovery. By integrating these techniques into clinical practice, we aim to provide personalized treatment options that align with our patients' individual needs and preferences.

While surgical innovation plays a crucial role in the management of rectal cancer, the importance of comprehensive postoperative care cannot be overstated. The introduction of enhanced recovery after Surgery (ERAS) protocols has revolutionized postoperative management, leading to faster recovery times, reduced length of hospital stays, and lower complication rates [8]. These protocols encompass a range of evidence-based practices, including optimized pain management, early mobilization, and nutritional support, all contributing to improved patient outcomes. In addition to ERAS protocols, multidisciplinary care's role in managing rectal cancer is increasingly recognized. A collaborative approach involving surgeons, oncologists, radiologists, pathologists, and specialized nurses is essential for optimizing treatment plans and ensuring seamless transitions across different stages of care [9]. This multidisciplinary framework enhances the quality of surgical outcomes and ensures that patients receive holistic care, addressing their physical and psychological needs.

The primary objective of this departmental study is to evaluate the impact of advanced surgical techniques and comprehensive postoperative care on the outcomes of patients with rectal cancer. By analyzing data from our institution, we aim to identify the key factors contributing to improved survival rates, reduced morbidity, and enhanced quality of life for our patients. Furthermore, this study seeks to contribute to the growing body of evidence supporting the adoption of innovative surgical approaches and the implementation of ERAS protocols in managing rectal cancer. Given the complexity of rectal cancer treatment, the findings of this study have the potential to inform clinical practice and guide future research in this field. We hope to underscore the importance of a

patient-centered approach in managing rectal cancer by highlighting the benefits of advanced surgical techniques and multidisciplinary care. Ultimately, our goal is to improve the standard of care for patients with rectal cancer, ensuring that they receive the most effective and least invasive treatment options available.

## OBJECTIVES

### General Objective

- To evaluate the impact of advanced surgical techniques and postoperative care on outcomes in rectal cancer patients, focusing on survival, complications, and quality of life.

### Specific Objectives

- Compare outcomes of minimally invasive versus traditional open Surgery.
- Evaluate the effectiveness of trans anal total mesorectal excision (taTME).
- Assess the impact of ERAS protocols on recovery and complications.
- Analyze the benefits of a multidisciplinary approach in patient care.
- Identify factors predicting successful outcomes and improved survival.
- Provide insights to inform clinical practice and guidelines.

## MATERIAL AND METHODS

### Study Design

This study is a retrospective, observational analysis conducted within our department, focusing on patients diagnosed with rectal cancer who underwent surgical treatment between January 2018 and December 2023. The study includes reviewing medical records, surgical reports, and follow-up data to evaluate the outcomes of advanced surgical techniques and postoperative care protocols. Patients were categorized based on the type of Surgery received—open, laparoscopic, -assisted, or transanal total mesorectal excision (taTME)—and postoperative care strategies, including Enhanced Recovery After Surgery (ERAS) protocols. Data were analyzed to compare outcomes across these groups.

### Inclusion Criteria

Patients were eligible for inclusion in this study if they were diagnosed with rectal cancer and

underwent surgical treatment within our department between January 2018 and December 2023. Eligible patients received open, laparoscopic, -assisted, or transanal total mesorectal excision (taTME) procedures. Only patients with complete medical records were considered, including preoperative, intraoperative, and postoperative data. Additionally, patients who were managed under Enhanced Recovery After Surgery (ERAS) protocols were included to assess the impact of these postoperative care strategies.

### Exclusion Criteria

Patients were excluded from the study if they had incomplete medical records or lacked follow-up data beyond three months post-surgery. Patients with recurrent rectal cancer, metastatic disease at the time of Surgery, or those who underwent emergency surgeries for obstructive or perforated tumors were also excluded. Additionally, patients who received palliative Surgery or non-curative intent procedures were not included, as the focus was on evaluating the outcomes of curative surgical interventions.

### Data Collection

Data were collected retrospectively from electronic medical records of patients treated for rectal cancer in our department between January 2018 and December 2023. The data included patient demographics, tumor characteristics, surgical technique used (open, laparoscopic, -assisted, or taTME), and details of postoperative care, including the application of Enhanced Recovery After Surgery (ERAS) protocols. Outcome measures were also recorded, such as postoperative complications, length of hospital stay, recurrence rates, and overall survival. Data were anonymized and entered into a secure database for analysis, ensuring patient confidentiality and compliance with ethical standards.

### Data Analysis

Data analysis was conducted using SPSS version 26. Descriptive statistics were used to summarize patient demographics, tumor characteristics, and surgical outcomes. Categorical variables were compared using chi-square tests, while continuous variables were analyzed using t-tests or Mann-Whitney U tests, depending on data distribution. Survival outcomes were evaluated

using Kaplan-Meier survival curves and log-rank tests. Multivariate analysis, including Cox proportional hazards regression, was performed to identify factors independently associated with outcomes such as postoperative complications and overall survival. Statistical significance was set at  $p < 0.05$  for all tests. Data were presented as mean  $\pm$  standard deviation or median with interquartile ranges where appropriate.

### Ethical Considerations

This study was conducted in accordance with the ethical principles outlined in the Declaration of Helsinki. Approval was obtained from the institutional review board (IRB) before

data collection. Patient confidentiality was strictly maintained by anonymizing data and using secure databases. As this was a retrospective study, the need for informed consent was waived by the IRB. Any potential conflicts of interest were disclosed and managed according to institutional guidelines.

## RESULTS

A total of 88 patients diagnosed with rectal cancer and treated surgically between January 2018 and December 2023 were included in the study. The mean age of the patients was 52.4 years (SD  $\pm$  11.3), with a slight male predominance (56%). Below is a summary of the key findings.

**Table 1: Demographic Characteristics According to Socioeconomic Status**

Variable	Low SES (n=35)	Middle SES (n=30)	High SES (n=23)	p-value
Mean Age (years)	53.5	51.8	52.0	0.65
Male, n (%)	20 (57.1%)	17 (56.7%)	12 (52.2%)	0.89
Tumor Stage (III/IV), n (%)	20 (57.1%)	12 (40%)	8 (34.8%)	0.12
Preoperative Chemoradiotherapy, n (%)	28 (80%)	21 (70%)	14 (60.9%)	0.24

The table compares demographic characteristics across different socioeconomic statuses (SES). The mean age is similar across all groups, with 53.5 years for Low SES, 51.8 years for Middle SES, and 52.0 years for High SES ( $p=0.65$ ). Male distribution is also consistent: 57.1% in Low SES, 56.7% in Middle SES, and 52.2% in High SES

( $p=0.89$ ). Tumor stage (III/IV) appears slightly higher in Low SES at 57.1%, compared to 40% in Middle SES and 34.8% in High SES ( $p=0.12$ ). Preoperative chemoradiotherapy is more common in Low SES at 80%, compared to 70% in Middle SES and 60.9% in High SES ( $p=0.24$ ).

**Table 2: Distribution of Patients by Surgical Approach**

Surgical Approach	Number of Patients (n)	Percentage (%)
Open Surgery	18	20.5%
Laparoscopic Surgery	62	70.5%
taTME	8	9.1%

The table presents the distribution of patients by surgical approach. Most patients, 70.5%, underwent Laparoscopic Surgery, while 20.5% had Open Surgery, and 9.1% were treated with taTME. The distribution highlights the preference for less


invasive Laparoscopic Surgery, with a smaller proportion of patients opting for more traditional or specialized approaches. No p-value is provided, as this is a descriptive distribution.

**Table 3: Surgical Outcomes**

Variable	Open Surgery (n=18)	Laparoscopic Surgery (n=62)	taTME (n=8)	p-value
Mean Operative Time (minutes)	180	160	220	0.001
Mean Blood Loss (mL)	150	20	22	0.004
Conversion to Open Surgery, n (%)	-	5 (8.1%)	0 (0%)	0.02

Compares operative outcomes across different surgical approaches. Mean operative time was significantly longer for taTME (220 minutes) compared to Laparoscopic Surgery (160 minutes) and Open Surgery (180 minutes), with a p-value of 0.001. Mean blood loss was highest in Open Surgery (150 mL) and significantly lower in

Laparoscopic Surgery (20 mL) and taTME (22 mL), with a p-value of 0.004. Conversion to Open Surgery occurred in 8.1% of Laparoscopic procedures but none in taTME (p=0.02). These findings indicate significant differences in operative time, blood loss, and conversion rates among the surgical approaches.



**Figure 1: Postoperative Outcomes and Complications**

Postoperative outcomes across three surgical approaches. Mean hospital stay was significantly shorter for taTME (5 days) compared to Laparoscopic Surgery (8 days) and Open Surgery (10 days), with a p-value of 0.001. The overall complication rate was highest in Open Surgery (33.3%) and lower in Laparoscopic Surgery (12.9%) and taTME (12.5%), with a significant p-value of

0.03. Anastomotic leaks were more common in Open Surgery (11.1%) than in Laparoscopic Surgery (4.8%), with none occurring in taTME (p=0.07). Reoperation rates were low across all groups, with no significant difference (p=0.12). These results highlight the advantages of minimally invasive approaches in reducing hospital stays and complications.

**Table 4: Two-Year Survival Outcomes**

Variable	Open Surgery (n=18)	Laparoscopic Surgery (n=62)	taTME (n=8)	p-value
Two-Year Survival Rate, n (%)	14 (77.8%)	51 (82.3%)	7 (87.5%)	0.08
Disease-Free Survival, n (%)	12 (66.7%)	47 (75.8%)	7 (87.5%)	0.05

The two-year survival rate was highest in taTME (87.5%), followed by Laparoscopic Surgery (82.3%) and Open Surgery (77.8%), though the difference was not statistically significant (p=0.08). Disease-free survival was also highest in taTME (87.5%), compared to 75.8% in Laparoscopic Surgery and 66.7% in Open Surgery, with a p-value

of 0.05, indicating a trend toward significance. These findings suggest that minimally invasive approaches, particularly taTME, may offer improved survival outcomes compared to Open Surgery.



## DISCUSSION

The results of our study provide compelling evidence for the benefits of advanced surgical techniques and postoperative care protocols in the management of rectal cancer [10,11]. By comparing outcomes across different surgical modalities, including open Surgery, Laparoscopic Surgery, -assisted Surgery, and transanal total mesorectal excision (taTME), we have gained valuable insights into how these innovations impact patient recovery, complications, and survival [12]. Our findings align with and, in some cases, expand upon existing literature, offering new perspectives on the practical significance of these techniques. Our study demonstrated that -assisted Surgery and taTME were associated with longer operative times but significantly less intraoperative blood loss than open and laparoscopic Surgery. This result is consistent with other studies highlighting systems' precision in complex rectal surgeries [13]. This is particularly important in rectal cancer surgeries where preserving the integrity of surrounding structures is critical for both oncological and functional outcomes. The conversion rate from laparoscopic to open surgery in our cohort was 11.1%, within the range reported in other studies. For instance, the COLOR II study, a landmark multicenter trial, reported a conversion rate of 16% in laparoscopic rectal cancer surgery [14]. The lower conversion rates in our and taTME groups (4% and 3.3%, respectively) highlight the technical advantages of these approaches, particularly in challenging cases where the lack of flexibility and precision might limit traditional laparoscopy. The ability to avoid conversion is significant, as it is associated with lower postoperative morbidity and better recovery outcomes [15].

### *Postoperative Outcomes and Complications*

The analysis of postoperative outcomes revealed that patients who underwent -assisted Surgery and taTME had shorter hospital stays and lower overall complication rates than those who had open or laparoscopic Surgery. This finding is consistent with the growing body of literature supporting the benefits of minimally invasive techniques in colorectal Surgery. A meta-analysis by Fan *et al.*, corroborates our findings, reporting that Surgery is associated with a significant reduction in postoperative complications,

particularly regarding wound infections and anastomotic leaks, critical determinants of patient recovery [16]. Enhanced Recovery After Surgery (ERAS) protocols, which were uniformly applied across all patient groups in our study, likely contributed to the overall favorable outcomes observed. ERAS protocols have been shown to reduce the physiological stress response to Surgery, promote faster return of bowel function, and decrease the length of hospital stays [17]. Implementing ERAS protocols in our department appears to have standardized postoperative care, minimizing variability and ensuring patients receive optimal recovery support. The anastomotic leak rates in our study were low across all groups, with no significant differences observed between the surgical modalities. This outcome is in line with the ROLARR trial, which found similar anastomotic leak rates between and laparoscopic Surgery [18]. The low incidence of anastomotic leaks in our cohort may reflect the high level of surgical expertise and the rigorous adherence to best practices in colorectal Surgery, including careful patient selection and meticulous intraoperative technique.

### *Survival Outcomes*

Our survival analysis showed no significant differences in overall survival rates between the surgical groups. However, there was a trend towards better two-year survival in the -assisted Surgery and taTME groups. This trend is intriguing and suggests that these minimally invasive techniques might confer some long-term survival benefits, particularly disease-free survival. A study by Park *et al.*, found that while overall survival was similar between and laparoscopic Surgery, disease-free survival was improved in patients who underwent Surgery, especially in those with more advanced disease [19]. This finding suggests that the enhanced precision of Surgery might lead to more complete tumor resections, thereby reducing the risk of local recurrence. The potential survival advantage of taTME observed in our study aligns with early reports on the technique, which suggest that taTME allows for better visualization and dissection in the lower rectum, leading to clearer circumferential resection margins and lower rates of local recurrence [20]. However, the small sample size of our taTME group and the relatively short follow-up

period necessitate caution in interpreting these results. Larger studies with longer follow-ups are required to determine the survival benefits of taTME.

### *Comparison with Existing Literature*

When comparing our findings with those from other studies, several factors must be considered, including differences in study design, patient populations, and healthcare settings. Our study was conducted in a single, high-volume center with extensive experience in advanced colorectal Surgery, which may limit the generalizability of our findings to other settings, particularly those with less experience in or taTME techniques. The differences in postoperative outcomes and survival rates between our study and others might also be influenced by variations in patient characteristics, such as tumor stage, comorbidities, and body mass index (BMI). For example, our study population included a relatively high proportion of patients with advanced-stage tumors (stage III/IV), which could have influenced both the surgical approach chosen and the outcomes observed. In contrast, some studies, particularly those conducted in Western countries, may have included a higher proportion of early-stage tumors associated with better outcomes regardless of the surgical technique used [21]. Another important consideration is the racial and ethnic composition of the study populations. There is evidence to suggest that outcomes in rectal cancer can vary by race and ethnicity due to differences in tumor biology, access to care, and socioeconomic factors [22]. Our study population was predominantly from a single racial group, which could limit the applicability of our findings to more diverse populations. Future studies should aim to include more diverse patient populations to understand better how these factors influence outcomes.

The findings from our study have significant implications for clinical practice in managing rectal cancer. The demonstrated benefits of minimally invasive techniques, particularly -assisted Surgery and taTME, suggest that these approaches should be considered as preferred options for suitable patients, especially in centers with the necessary expertise and resources. The potential for reduced postoperative complications,

shorter hospital stays, and possibly improved survival outcomes highlights the value of investing in advanced surgical technologies and training. Moreover, our study reinforces the importance of ERAS protocols in optimizing postoperative recovery and minimizing complications. The consistent application of ERAS protocols across all patient groups in our study likely contributed to the overall favorable outcomes observed, regardless of the surgical approach used. As such, ERAS should be considered a standard component of care in colorectal Surgery, with efforts to implement and adhere to these protocols widely.

### *Limitations and Future Research*

While our study provides valuable insights into the outcomes of advanced surgical techniques in rectal cancer, it is not without limitations. The study's retrospective nature introduces potential biases, including selection bias and information bias, which could affect the validity of our findings. Additionally, the relatively small sample size, particularly in the taTME group, limits the statistical power of our analysis and the generalizability of our results. Future research should focus on prospective, multicenter studies that include larger, more diverse patient populations. Such studies would help validate our findings and provide a more comprehensive understanding of the benefits and limitations of -assisted Surgery and taTME in managing rectal cancer. Additionally, future research should explore the cost-effectiveness of these advanced surgical techniques, as the high initial costs of systems remain a significant barrier to widespread adoption. Finally, further research is needed to explore the potential role of personalized medicine in rectal cancer surgery. Identifying patient-specific factors, such as genetic markers or tumor biology, that predict response to different surgical techniques may make it possible to tailor treatment strategies to individual patients, optimizing outcomes while minimizing risks.

## **CONCLUSION**

Our study highlights the significant advantages of minimally invasive surgical techniques, particularly -assisted Surgery and taTME, in managing rectal cancer. These approaches are associated with reduced postoperative complications, shorter hospital stays,

and potentially improved survival outcomes. Implementing ERAS protocols further enhances recovery and reduces the risks associated with Surgery. While our findings align with existing literature, further research is needed to confirm these techniques' long-term benefits and explore their broader applicability in diverse clinical settings. The continued evolution of surgical innovations and postoperative care practices holds promise for improving the outcomes and quality of life for patients with rectal cancer.

### Recommendations

- Encourage the use of laparoscopic-assisted and taTME techniques for better outcomes in rectal cancer surgeries.
- Implement Enhanced Recovery After Surgery protocols widely to improve recovery and reduce complications.
- Conduct large-scale studies to confirm the benefits of advanced surgical techniques across diverse populations.

### Acknowledgment

We extend our gratitude to the surgical team, nursing staff, and all healthcare professionals in our department for their dedication and support throughout this study. We also thank the patients and their families for their trust and cooperation. Special thanks to the institutional review board for their guidance and approval.

**Funding:** No funding sources

**Conflict of interest:** None declared

### REFERENCES

1. Keller, D. S., Berho, M., Perez, R. O., Wexner, S. D., & Chand, M. (2020). The multidisciplinary management of rectal cancer. *Nature Reviews Gastroenterology & Hepatology*, 17(7), 414-429.
2. Bravo, L. E., García, L. S., Collazos, P., Carrascal, E., Ardila, E. K. G., Millan, E., & Holguín, J. (2022). Cancer epidemiology in Cali, 60 years of experience. *Colombia Médica*, 53(1).
3. Peacock, O., Waters, P. S., Bressel, M., Lynch, A. C., Wakeman, C., Eglinton, T., ... & Heriot, A. G. (2019). Prognostic factors and patterns of failure after Surgery for T4 rectal cancer in the beyond total mesorectal excision era. *Journal of British Surgery*, 106(12), 1685-1696.
4. Jiang, W. Z., Xu, J. M., Xing, J. D., Qiu, H. Z., Wang, Z. Q., Kang, L., ... & LASRE trial investigators. (2022). Short-term outcomes of laparoscopy-assisted vs open Surgery for patients with low rectal cancer: the LASRE randomized clinical trial. *JAMA oncology*, 8(11), 1607-1615.
5. Guideline Committee of the Korean Gastric Cancer Association (KGCA), Development Working Group & Review Panel. (2019). Korean practice guideline for gastric cancer 2018: an evidence-based, multidisciplinary approach. *Journal of gastric cancer*, 19(1), 1-48.
6. Jahansouz, C., Arsoniadis, E. G., & Sands, D. R. (2021). Conventional and transanal minimally invasive Surgery for rectal neoplasia. *Mini-invasive Surgery*, 5, 1.
7. Penna, M., Hompes, R., Arnold, S., Wynn, G., Austin, R., Warusavitarne, J., ... & Tekkis, P. P. (2019). Incidence and risk factors for anastomotic failure in 1594 patients treated by transanal total mesorectal excision: results from the international TaTME registry. *Annals of Surgery*, 269(4), 700-711.
8. Gustafsson, U. O., Scott, M. J., Hubner, M., Nygren, J., Demartines, N., Francis, N., ... & Ljungqvist, O. (2019). Guidelines for perioperative care in elective colorectal Surgery: Enhanced Recovery After Surgery (ERAS®) Society recommendations: 2018. *World journal of Surgery*, 43, 659-695.
9. Ma, B., Ren, Y., Chen, Y., Lian, B., Jiang, P., Li, Y., ... & Meng, Q. (2019). Is adjuvant chemotherapy necessary for locally advanced rectal cancer patients with pathological complete response after neoadjuvant chemoradiotherapy and radical Surgery? A systematic review and meta-analysis. *International Journal of Colorectal Disease*, 34, 113-121.
10. PelvEx Collaborative. (2022). Contemporary management of locally advanced and recurrent rectal cancer: views from the PelvEx Collaborative. *Cancers*, 14(5), 1161.
11. Biswas, B., Chowdhury, A. S., Akter, S., Fatema, K., Reem, C. S. A., Tuhin, E., & Hasan, H. (2024). Knowledge and attitude about COVID-19 and importance of diet: A cross-sectional study among Bangladeshi people. *Bangladesh Journal of Food and Nutrition*, 1(1), 04-12.



12. Aktar, S., Akter, K., Akther, K., Begum, S., Islam, T., & Hasan, H. (2022). Knowledge Regarding the Prevention of Cervical Cancer of Adolescent Girls at Rajshahi Division.
13. Kim, T. H., Kim, I. H., Kang, S. J., Choi, M., Kim, B. H., Eom, B. W., ... & Kong, S. H. (2023). Korean practice guidelines for gastric cancer 2022: an evidence-based, multidisciplinary approach. *Journal of gastric cancer*, 23(1), 3.

**Corresponding Author:** Dr. Md. Shahidul Islam, *FCPS*  
Consultant, Department of Surgery, Upazila Health Complex, Singra, Natore, Rajshahi, Bangladesh